

Global Wild Poliovirus 2015 - 2020



Country or territory ³	Wild virus type 1 confirmed cases							Wild virus type 1 reported from other sources ²								
	Full year total					01 Jan -09 Jun ¹		Date of most recent case	Full year total					01 Jan -09 Jun ¹		Date of most recent virus
	2015	2016	2017	2018	2019	2019	2020		2015	2016	2017	2018	2019	2019	2020	
Pakistan	54	20	8	12	147	22	50	11-May-20	84	62	110	141	391	140	187	14-May-20
Afghanistan	20	13	14	21	29	8	12	22-Mar-20	20	2	42	83	56	23	13	27-Apr-20
Nigeria ³	0	4	0	0	0	0	0	21-Aug-16		1 ⁴						27-Sep-16
Iran	0	0	0	0	0	0	0	NA					3			20-May-19
Total (Type1)	74	37	22	33	176	30	62		104	64	65	224	450	163	200	
Tot. in endemic countries	74	37	22	33	176	30	62									
Tot. in non-end countries	0	0	0	0	0	0	0									
No. of countries (infected)	2	3	2	2	2	2	2									
No. of countries (endemic)	3	2 ³	2 ³	3	3	3	3									
Total Female	30	13	7	18	72	14	28									
Total Male	44	24	15	15	104	16	34									

Countries in yellow are endemic.¹Data reported to WHO HQ on 10 June 2019 for 2019 data and 09 June 2020 for 2020 data.

²Wild viruses from environmental samples, selected contacts, healthy children and other sources. Last WPV type 3 had its onset on 10 November 2012. ³Between 27 Sep 2015 and 27 Sep 2016, Nigeria was not classified as endemic. NA - Most recent case had onset prior to 1999. ⁴Exceptionally reporting case-contact of a positive index case given the date of collection is later than the onset date of the most recent WPV.